

## **Coding & Documentation**

The presence of a condition cannot be assumed even if evidence is present in the medical record. It requires the provider's clinical judgment and explicit diagnosis.

**Substance use disorder:** Patterned use of a substance in an amount or method causes significant impairment or distress.

**Physical Dependence:** A substance is needed to function normally; withdrawal symptoms would occur if the drug were stopped abruptly.

**Psychological Dependence:** Discontinued use of a substance results in emotional or behavioral disturbances.

#### Diagnostic Criteria<sup>1</sup>

Patient must display two or more of the following symptoms in a 12-month period:

- Use of substance in larger doses or for a longer time period than intended
- Persistent desire or unsuccessful efforts to reduce or control use
- Abundance of time spent obtaining, using or recovering from the effects of a substance
- Strong cravings or desire to use

- Recurrent use resulting in failure to fulfill obligations at work, home or school
- Continued use even when persistent or recurrent social or interpersonal problems are caused by the effects
- Abandonment of important social, recreational or occupational activities because of use
- Recurrent use in physically hazardous situations
- Tolerance the need for increased amount achieve desired effect; the diminished effect with continued use of same amount
- Withdrawal characteristic withdrawal syndrome; use to relieve or ovoid withdrawal

#### ICD-10-CM Codes<sup>2</sup>

Mental and behavioral disorders due to psychoactive substance use, F10-F19

**Identify the substance** — each substance is addressed as a separate disorder:

- Alcohol
- Opioid
- Cannabis
- Sedative, hypnotic or anxiolytic

- Cocaine
- Other stimulant (amphetamine, caffeine, etc.)
- Hallucinogen
- Nicotine

- Inhalant
- Other (or unknown) psychoactive substance

The educational material herein complies with accepted ICD-10 guidelines and is for general supplemental purposes only. The information herein is not guaranteed to be complete, free of errors, or the most current revision. It is the responsibility of the provider to document accurate and complete codes, clinical rationale, and medical services rendered to support appropriate ICD-10 code(s) according to official billing and coding guidelines, procedures, and regulations.

<sup>1</sup> American Psychiatric Association: Diagnostic and Statistical Manual of Mental Disorders: Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition. Arlington, VA: American Psychiatric Association, 2013.

<sup>2</sup> https://www.cms.gov/medicare/icd-10/2021-icd-10-cm



### **Coding Hierarchy**

If Documented	Then Code
Use & abuse	Abuse
Use & dependence	Dependence
Abuse & dependence	Dependence
Use, abuse & dependence	Dependence

### **Disorder Severity**

Mild	2–3 symptoms
Moderate	4-5 symptoms
Severe	6+ symptoms

#### Severe use disorder = addiction3

A substance addiction will likely require ongoing treatment for several years.

#### "In Remission"

Disorder	Classification
Early remission	No signs/symptoms of disorder for 3–12 months
Sustained remission	No signs/symptoms of disorder for 12+ months
Abuse in remission	Mild use disorder in early or sustained remission
Dependence in remission	Moderate or severe use disorder in early or sustained remission

### Document to the highest known degree and code to the highest specificity.

Documentation should include, if applicable:

- Substance type
- Severity
- Association with intoxication or withdrawal
- Detailed clinical diagnosis
- Treatment plan/condition management

- Presence of induced manifestation or complication, such as:
  - Delirium
  - Perceptual disturbance
  - Anxiety
  - Sexual dysfunction
  - Sleep or mood disorder
  - Persisting amnestic disorder or dementia

<sup>3</sup> https://www.addictionpolicy.org/post/dsm-5-facts-and-figures



## **HEDIS® & Quality**

### Initiation and Engagement of Alcohol & Other Drug Abuse or Dependence Treatment (IET)

Measure evaluates adolescent and adult members, ages 13 and older, with a new episode of alcohol or other drug (AOD) abuse or dependence who received the following between January 1 and November 14 of the measurement year

#### Two rates are reported:

Initiation of AOD Treatment	Engagement of AOD Treatment
intensive outpatient encounter or partial hospitalization, telehealth, or medication	Members who initiated treatment and were engaged in ongoing AOD treatment within 34 days of the initiation visit. Engagement is at least one medication treatment event or two visits by AOD admission, outpatient visit, intensive outpatient encounter, or partial hospitalization, or telehealth.

#### Initiation of AOD Treatment

Submit claims with any diagnosis code to identify AOD with an IET treatment code as listed below.

#### For all initiation events except medication treatment, initiation on the same day as the index episode start date (IESD) must be with different providers in order to count.

Outpatient, intensive outpatient, partial hospitalization, observation, telehealth, or ED visit (not resulting in an inpatient stay)	The date of service
Inpatient stay or for detoxification that occurred during an inpatient stay	The date of discharge
Detoxification (other than detoxification that occurred during an inpatient stay)	The date of service
ED or observation visits that result in an inpatient stay	The date of the inpatient discharge
Direct transfers	The discharge date from the last admission

For additional resources, contact our provider relations team at Providers@ARHealthWellness.com



- If the index episode was an inpatient discharge or ED/observation visit that resulted in an inpatient stay, then the inpatient stay is considered initiation of treatment and the member is compliant.
- . If the index episode was an opioid treatment service that bills monthly, then the opioid treatment service is considered initiation of treatment and the member is compliant: HCPCS codes G2086, G2087
- If the index episode was not an inpatient discharge, then the member must initiate treatment on the IESD or in the 13 days following the IESD (14 total days). Any of the following combinations must meet criteria for initiation.

ICD-10-CM codes for alcohol, opioid, or other drug abuse or dependence are listed on the next page.

Description	Coding			
An acute or non-acute inpatient	UBREV			
admission <b>with</b> an alcohol, opioid or other drug abuse and/ or dependence diagnosis on the discharge claim	0100, 0101, 0110, 0111, 0112, 0113, 0114, 0116, 0117, 0118, 0119, 0120, 0121, 0122, 0123, 0124, 0126, 0127, 0128, 0129, 0130, 0131, 0132, 0133, 0134, 0136, 0137, 0138, 0139, 0140, 0141, 0142, 0143, 0144, 0146, 0147, 0148, 0149, 0150, 0151, 0152, 0153, 0154, 0156, 0157, 0158, 0159, 0160, 0164, 0167, 0169, 0170, 0171, 0172, 0173, 0174, 0179, 0190, 0191, 0192, 0193, 0194, 0199, 0200, 0201, 0202, 0203, 0204, 0206, 0207, 0208, 0209, 0211, 0212, 0213, 0214, 0219, 1000, 1001, 1002			
	CPT®	HCPCS		UBREV
IET standalone visits <b>with</b> an alcohol, opioid or other drug abuse and/or dependence diagnosis	98960, 98961, 98962, 99078, 99201, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99241, 99242, 99243, 99244, 99245, 99341, 99342, 99343, 99344, 99345, 99347, 99348, 99349, 99350, 99384, 99385, 99386, 99387, 99394, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99408, 99409, 99411, 99412, 99483, 99510	G0410, G0411, G0443 H0004, H0005, H00 H0031, H0034, H003 H0040, H0047, H200 H2012, H2013, H2014,	G0396, G0397, G0409, G0463, H0001, H0002, 07, H0015, H0016, H0022, 5, H0036, H0037, H0039, 00, H2001, H2010, H2011, H2015, H2016, H2017, 0, H2035, H2036, S0201, 1, T1006, T1012, T1015	0510, 0513, 0515, 0516, 0517, 0519, 0520, 0521, 0522, 0523, 0526, 0527, 0528, 0529, 0900, 0902, 0903, 0904, 0905, 0906, 0907, 0911, 0912, 0913, 0914, 0915, 0916, 0917, 0919, 0944, 0945, 0982, 0983
An observation <b>with</b> an alcohol,	СРТ			
opioid or other drug abuse and/ or dependence diagnosis 99217, 99218, 99219, 99220				
an alcohol, opioid or other drug abuse and/or dependence	СРТ		Place of Service	
	90791, 90792, 90832, 90833, 90834, 90836, 90837, 90838, 90839, 90840, 90845, 90847, 90849, 90853, 90875, 90876		02, 03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 52, 53, 57, 58, 71, 72	



Description	Coding		
IET group 2 visits with POS and	СРТ		Place of Service
an alcohol, opioid or other drug abuse and/or dependence diagnosis	99221, 99222, 99223, 99231, 99232, 99233, 99238, 99239, 99251, 99252, 99253, 99254, 99255		02, 52, 53
A telephone visit <b>with</b> an alcohol,	СРТ		
opioid or other drug abuse and/ or dependence diagnosis	98966, 98967, 98968, 99441, 99442, 99443		
An e-visit or virtual check-in	СРТ	HCPCS	
with an alcohol, opioid or other drug abuse and/ or dependence diagnosis	98969, 98970, 98971, 98972, 99421, 99422, 99423, 99444, 99458	G2010, G20	112, G2061, G2062, G2063

#### Alcohol, Opioid, and Other Drug Abuse or Dependence ICD-10-CM Codes

F10.10, F10.120, F10.121, F10.129, F10.14, F10.150, F10.151, F10.159, F10.180, F10.181, F10.182, F10.188, F10.19, F10.20, F10.220, F10.221, F10.229, F10.230, F10.231, F10.232, F10.232, F10.239, F10.24, F10.250, F10.251, F10.259, F10.26, F10.27, F10.280, F10.281, F10.282, F10.288, F10.29, F11.10, F11.121, F11.122, F11.129, F11.14, F11.150, F11.151, F11.159, F11.181, F11.182, F11.188, F11.19, F11.20, F11.221, F11.222, F11.229, F11.23, F11.24, F11.250, F11.251, F11.259, F11.281, F11.282, F11.288, F11.29, F12.10, F12.120, F12.121, F12.122, F12.129, F12.150, F12.151, F12.159, F12.180, F12.189, F12.19, F12.20, F12.220, F12.221, F12.222, F12.229, F12.23, F12.250, F12.251, F12.259, F12.280, F12.288, F12.29, F13.10, F13.120, F13.121, F13.129, F13.14, F13.150, F13.151, F13.159, F13.181, F13.182, F13.188, F13.19, F13.20, F13.220, F13.221, F13.229, F13.230, F13.231, F13.232, F13.239, F13.24, F13.250, F13.251, F13.259, F13.26, F13.27, F13.280, F13.281, F13.282, F13.288, F13.29, F14.10, F14.120, F14.121, F14.122, F14.129, F14.14, F14.150, F14.151, F14.159, F14.180, F14.181, F14.182, F14.188, F14.19, F14.20, F14.220, F14.221, F14.222, F14.229, F14.23, F14.244, F14.250, F14.251, F14.259, F14.280, F14.281, F14.282, F14.288, F14.29, F15.10, F15.120, F15.121, F15.122, F15.129, F15.14, F15.150, F15.151, F15.159, F15.180, F15.181, F15.182, F15.188, F15.19, F15.200, F15.220, F15.221, F15.222, F15.222, F15.222, F15.222, F15.223, F15.234, F15.250, F15.251, F15.259, F15.280, F15.281, F15.282, F15.288, F15.29, F16.10, F16.120, F16.121, F16.122, F16.129, F16.14, F16.150, F16.151, F16.159, F16.180, F16.183, F16.19, F16.20, F16.220, F16.221, F16.229, F16.244, F16.250, F16.251, F16.259, F16.280, F16.283, F16.288, F18.29, F18.10, F18.121, F18.129, F18.14, F18.150, F18.151, F18.159, F18.10, F18.180, F18.200, F18.221, F18.229, F18.24, F18.250, F18.250, F19.220, F19.221, F19.222, F19.222, F19.222, F19.222, F19.222, F19.222, F19.222, F19.222, F19.2



If the index episode was for a diagnosis of alcohol abuse or dependence, a medication treatment-dispensing event or medication treatment during a visit, then:

HCPCS Codes	ICD-10-CM Codes
H0020, H0033, J0570, J0571, J0572, J0573, J0574, J0575, J2315, Q9991, Q9992, S0109	F10.10, F10.120, F10.121, F10.129, F10.14, F10.150, F10.151, F10.159, F10.180, F10.181, F10.182, F10.188, F10.19, F10.20, F10.220, F10.221, F10.229, F10.230, F10.231, F10.232, F10.239, F10.24, F10.250, F10.251, F10.259, F10.26, F10.27, F10.280, F10.281, F10.282, F10.288, F10.29

Alcohol Use Disorder Treatment Medications		
Description	Prescription	
Aldehyde dehydrogenase inhibitor	Disulfiram (oral)	
Antagonist	Naltrexone (oral and injectable)	
Other	Acamprosate (oral; delayed-release tablet)	

If the index episode was for a diagnosis of opioid abuse or dependence during a visit, then:

HCPCS Codes	ICD-10-CM Codes
G2071, G2074, G2075, G2076, G2077, G2080, G086, G087	F11.10, F11.120, F11.121, F11.122, F11.129, F11.14, F11.150, F11.151, F11.159, F11.181, F11.182, F11.188, F11.19, F11.20, F11.220, F11.221, F11.222, F11.229, F11.23, F11.24, F11.250, F11.251, F11.259, F11.281, F11.282, F11.288, F11.29

If the index episode was for a diagnosis of opioid abuse or dependence, a medication treatment-dispensing event or medication treatment during a visit, then:

HCPCS Codes	ICD-10-CM Codes
	F11.10, F11.120, F11.121, F11.122, F11.129, F11.14,
H0020, H0033, J0570, J0571, J0572,	F11.150, F11.151, F11.159, F11.181, F11.182,
J0573, J0574, J0575, J2315, Q9991, Q9992,	F11.188, F11.19, F11.20, F11.220, F11.221,
S0109, G2067, G2068, G2069 ,G2070,	F11.222, F11.229, F11.23, F11.24, F11.250,
G2072, G2073	F11.251, F11.259, F11.281, F11.282, F11.288,
	F11.29

Alcohol Use Disorder Treatment Medications		
Description	Prescription	
Antagonist	Naltrexone (oral and injectable)	
Partial agonist	<ul> <li>Buprenorphine (sublingual tablet, injection, implant)</li> <li>Buprenorphine/naloxone (sublingual tablet, buccal film, sublingual film)</li> </ul>	



### **Engagement of AOD Treatment**

Submit claims with any diagnosis code to identify AOD with an IET treatment code as listed below:

- Two engagement visits can be on the same date of service, but they must be with different providers in order to count as two events. An engagement visit on the same date of service as an engagement medication treatment event meets criteria, and there is no requirement that they be with different providers.
- For members who initiated treatment via an inpatient admission, the 34-day period for engagement begins the day after discharge.
- If a member had an opioid treatment service that bills monthly, or if they had a visit that included medication administration beginning the day after the initiation encounter through 34 days following the initiation event, and if the IESD was a diagnosis of opioid abuse or dependence, then the member is compliant with engagement for AOD treatment.
- If a member's initiation of AOD treatment was a medication treatment event, and the member has two or more engagement events in which only one can be an engagement medication treatment event, beginning the day after the initiation encounter through 34 days following the initiation event (34 total days), then the member is compliant with engagement for AOD treatment: HCPCS codes G2086, G2087, G2068, G2069, G2070, G2072, G2073

For additional resources, contact our provider relations team at Providers@ARHealthWellness.com



Any of the following beginning the day after the initiation encounter through 34 days following the initiation event (34 total days) meet criteria for an engagement visit:

Description	Coding				
	UBREV				
An acute or non-acute inpatient admission <b>with</b> an alcohol, opioid, or other drug abuse and/or dependence diagnosis on the discharge claim	0100, 0101, 0110, 0111, 0112, 0113, 0114, 0116, 0117, 0118, 0119, 0120, 0121, 0122, 0123, 0124, 0126, 0127, 0128, 0129, 0130, 0131, 0132, 0133,				
	СРТ	HCPCS		UBREV	
IET standalone visits <b>with</b> an alcohol, opioid, or other drug abuse and/or dependence diagnosis	98960, 98961, 98962, 99078, 99201, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99241, 99242, 99243, 99244, 99245, 99341, 99342, 99343, 99344, 99345, 99347, 99348, 99349, 99350, 99384, 99385, 99386, 99387, 99394, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99408, 99409, 99411, 99412, 99483, 99510	G0410, G0411, G0443 H0004, H0005, H00 H0031, H0034, H003 H0040, H0047, H200 H2012, H2013, H2014	G0396, G0397, G0409, 8, G0463, H0001, H0002, 07, H0015, H0016, H0022, 55, H0036, H0037, H0039, 00, H2001, H2010, H2011, 1, H2015, H2016, H2017, 10, H2035, H2036, S0201, 15, T1006, T1012, T1015	0510, 0513, 0515, 0516, 0517, 0519, 0520, 0521, 0522, 0523, 0526, 0527, 0528, 0529, 0900, 0902, 0903, 0904, 0905, 0906, 0907, 0911, 0912, 0913, 0914, 0915, 0916, 0917, 0919, 0944, 0945, 0982, 0983	
An observation <b>with</b> an alcohol,	СРТ				
opioid, or other drug abuse and/or dependence diagnosis	99217, 99218, 99219, 99220				
IET group 1 visits <b>with</b> POS <b>with</b> an alcohol, opioid, or other drug abuse and/or dependence diagnosis	CPT Place of		Place of Service	e of Service	
	90791, 90792, 90832, 90833, 90834, 90836, 90837, 90838, 90839, 90840, 90845, 90847, 90849, 90853, 90875, 90876		02, 03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 52, 53, 57, 58, 71, 72		
IET group 2 visits <b>with</b> POS <b>with</b> an alcohol, opioid, or other drug abuse and/or dependence diagnosis	СРТ		Place of Service		
	99221, 99222, 99223, 99231, 99232, 99233, 99238, 99239, 99251, 99252, 99253, 99254, 99255		02, 52, 53		
A telephone visit <b>with</b> an alcohol,	СРТ				
opioid or other druge abuse and/or dependence diagnosis	98966, 98967, 98968, 99441, 99442, 99443				
An e-visit or virtual check-in <b>with</b> an alcohol, opioid, or other drug abuse and/or dependence diagnosis	CPT HCPCS				
	98969, 98970, 98971, 98972, 99421, 99422, 99423, 99444, 99458		G2010, G2012, G2061, G2062, G2063		



If the index episode was for a diagnosis of opioid abuse or dependence during an opioid treatment service, then:

HCPCS Codes	ICD-10-CM Codes
2071, G2074, G2075, G2076, G2077, G2080, G086, G087 F11.10, F11.120, F11.121, F11.122, F11.129, F11.14, F11.150, F11.151, F11.159, F11.187 F11.19, F11.20, F11.20, F11.220, F11.221, F11.229, F11.229, F11.23, F11.24, F11.250, F11.	
	F11.281, F11.282, F11.288, F11.29

If the IESD diagnosis was a diagnosis of alcohol abuse or dependence, one or more medication treatment-dispensing events, or medication treatment during a visit, then beginning the day after the initiation encounter through 34 days following the initiation event (34 total days), this meets criteria for alcohol abuse and dependence treatment.

HCPCS Codes	ICD-10-CM Codes
H0020, H0033, J0570, J0571, J0572, J0573, J0574, J0575, J2315, Q9991, Q9992, S0109	F10.10, F10.120, F10.121, F10.129, F10.14, F10.150, F10.151, F10.159, F10.180, F10.181, F10.182, F10.188, F10.19, F10.20, F10.220, F10.221, F10.229, F10.230, F10.231, F10.232, F10.239, F10.24, F10.250, F10.251, F10.259, F10.26, F10.27, F10.280, F10.281, F10.282,
	F10.288, F10.29

Alcohol Use Disorder Treatment Medications		
Description	Prescription	
Aldehyde dehydrogenase inhibitor	Disulfiram (oral)	
Antagonist	Naltrexone (oral and injectable)	
Other	Acamprosate (oral; delayed-release tablet)	

If the IESD was a diagnosis of opioid abuse or dependence, one or more medication-dispensing events, or medication treatment during a visit, then beginning the day after the initiation encounter through 34 days following the initiation event (34 total days), this meets criteria for opioid abuse and dependence treatment.

HCPCS Codes	ICD-10-CM Codes
H0020, H0033, J0570, J0571, J0572,	F11.10, F11.120, F11.121, F11.122, F11.129, F11.14,
J0573, J0574, J0575, J2315, Q9991, Q9992,	F11.150, F11.151, F11.159, F11.181, F11.182,
S0109	F11.188, F11.19, F11.20, F11.220, F11.221,
	F11.222, F11.229, F11.23, F11.24, F11.250,
	F11.251, F11.259, F11.281, F11.282, F11.288,
	F11.29

Opioid Use Disorder Treatment Medications		
Description	Prescription	
Antagonist	Naltrexone (oral and injectable)	
Partial agonist	<ul> <li>Buprenorphine (sublingual tablet, injection, implant)</li> <li>Buprenorphine/naloxone (sublingual tablet, buccal film, sublingual film)</li> </ul>	